

Instructions Following Quadriceps or Patellar Tendon Repair

1. When to contact Dr. Faulkner's team:

- Excessive bleeding; unrelenting pain; excessive wound drainage (mild drainage or spotting is normal for 3-4 days postop); persistent numbness or tingling in operative extremity; temperature of $>101^{\circ}$ F; increasing redness or swelling; calf pain, tenderness, or swelling

2. Dressings/wound care:

- A rectangular waterproof dressing will cover your incision. Leave in place until 1st postop appointment.
- If your dressing happens to be removed, cover with gauze and paper tape.
- Ok to shower 2 days postop. Ok to let soapy water run over dressing. Do not submerge your incision.

3. Blood Clot Prevention:

- Wear compression stockings for 14 days postop. Remove for hygiene and for 2 hours/day to allow the skin to breathe.
- Take Aspirin 325 mg for 14 days.

4. Brace:

- Wear locked in full extension at all times, including at night.
- Unlock at least 3x/day for knee range of motion (0-30 degrees until further instructed).
- Ok to remove for hygiene.

5. Weight Bearing: Ok to place full weight on operative leg immediately after surgery with the knee brace locked in extension and using crutches. Wean off the crutches as tolerated.

6. Activity after surgery:

- Elevate leg above heart level as much as possible to help reduce swelling.
- Start knee range of motion exercises the day after surgery (Goal is 0-30° of flexion by the 1st postop visit).
- Quad sets and straight leg raises should be performed at least 3x daily.

7. Restrictions/Driving: No driving unless cleared by Dr. Faulkner. No lifting, pushing, pulling, or carrying objects > 10 lbs. Do not climb, stoop, crawl, or kneel.

8. Cold therapy/Ice:

- Apply an ice pack or cold therapy unit to knee 30 minutes 6 x daily to help reduce pain and swelling.
 - Do not apply to bare skin. Be sure to have an ACE wrap or wash cloth between ice pack and skin to avoid frostbite.
- Remove for 30 minutes every 2 hours.



NATHAN FAULKNER, MD

ORTHOPEDIC SURGERY • SPORTS MEDICINE

9. Pain medication:

- Take 1-2 tabs of the prescribed pain medicine with food every 4-6 hours as needed. Take the medication sparingly and discontinue as soon as possible.
- Common side effects of narcotics: constipation, drowsiness, nausea and confusion. If you experience severe nausea, itching, dizziness or other adverse reactions, call Dr. Faulkner.
- Increase your fiber intake (fruits, vegetable, grains) and use an over the counter stool softener (such as docusate or Senna-S) if you are having constipation.
- NOTE: New prescriptions for narcotic pain medication cannot be called in.
- Substitute Tylenol and/or Ibuprofen or Aleve if you do not have any contraindication to taking these. **Maximum dose/day of Tylenol is 1000 mg every 8 hours and Ibuprofen 800 mg every 8 hours.** Stomach upset, GI ulcers and allergic reactions are the most common adverse reactions to anti-inflammatories. You should not take more than 3000 mg of Tylenol in a 24 hour period.

10. Diet: Anesthesia can cause temporary nausea after surgery. Begin with light with clear liquids or soft foods after surgery to minimize upset stomach. Drink at least 6-8 glasses of water daily.

Website URL: nathanfaulknermd.com

Postop instruction videos URL: nathanfaulknermd.com/postop

Videos to watch:

1. *Post Op: Get to Know Your Knee Brace*
2. *Post Op: Knee Home Exercise Program*
3. *Shoulder & Knee Arthroscopy Post Op Wound Care Instructions*



SCAN ME

Please feel free to contact our office at any time!



NATHAN FAULKNER, MD

ORTHOPEDIC SURGERY • SPORTS MEDICINE